



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Diffusive isotope fractionation in silicate liquids: Dependence on liquid composition, cation bonding, and isotopic exchange

J. M. Watkins, D. J. DePaolo, F. J. Ryerson

September 14, 2010

AGU Fall Meeting 2010
San Francisco, CA, United States
December 13, 2010 through December 17, 2010

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

Diffusive isotope fractionation in silicate liquids: Dependence on liquid composition, cation bonding, and isotopic exchange

James M. Watkins, Donald J. DePaolo, and Frederick J. Ryerson

Molecular diffusion in natural volcanic liquids discriminates between isotopes of major elements. Although isotope separation by diffusion is expected on theoretical grounds, the dependence on mass is highly variable for different elements and in different media. We have been using diffusion couple experiments to measure the relative diffusivities of Ca and Mg isotopes in both natural and synthetic silicate liquids of varying composition. Our recent experimental results, in combination with earlier results from natural volcanic liquids, show clearly that the efficiency of diffusive separation of Ca isotopes varies considerably with liquid composition, yet is systematically related to the normalized diffusivity - the ratio of the mobility of the cation (D_{Ca}) to the mobility of the liquid matrix (D_{Si}). The results on Ca isotopes are consistent with available data on Fe, Li, and Mg isotopes in silicate liquids, when considered in terms of the parameter D_{cation}/D_{Si} .

Large D_{cation}/D_{Si} corresponds to a greater efficiency of isotope separation during diffusion; in this regime, we infer that the cation is loosely bound to larger aluminosilicate structures in the liquid. As D_{cation}/D_{Si} approaches unity, the efficiency of isotope separation is reduced, presumably because the cation is more strongly bound to the silicate liquid matrix. Our experiments on natural volcanic liquids demonstrate that diffusive coupling between Ca and Al (or Si) can lead to large $\delta^{44}\text{Ca}$ variations, even where there are small gradients in Ca concentration. These effects combined with theoretical models of diffusion allow us to move closer to a general model of isotopic fractionation during diffusion in silicate liquids, and to shed light on the bonding of cations within the liquids and the structure of the liquids themselves.

All of the experiments done thus far show diffusive isotope effects that depart from those predicted by an effective binary diffusion model, where each isotope diffuses in response to its own chemical gradient. Experiments with large D_{cation}/D_{Si} show only slight misfits, which can be explained by isotopic exchange superimposed on chemical diffusion. We propose a new model that accounts for this effect, in which the diffusivity describing the rate of isotope redistribution (D_s) is greater than that describing chemical diffusion (D_c) by a factor of 2 to 6. The new model provides a simple explanation for what has been described as chemical and isotopic “decoupling” during diffusion in silicate liquids. In natural liquids where D_{cation}/D_{Si} is close to unity, much larger deviations from simple diffusion models are observed. We are exploring alternative models of diffusion and the role that non-ideal mixing may play in generating these unexpected isotopic effects.

This work performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.